

100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

N

[illegible]

NOVEMBER, 1899

## SOME REMARKS ON THE SYMPTOMS OF BRONCHOCELE AND THE RESULTS OF OPERATIVE TREATMENT

BY FRANCIS J. SHEPHERD, M.D.,  
OF MONTREAL.

Professor of Anatomy and Lecturer on Operative Surgery, McGill University; Senior Surgeon to the Montreal General Hospital.

SC

ma  
all  
sid  
thi  
con  
be  
flu  
ria  
rha  
In  
the  
chy  
esp  
a f  
per  
ber  
ioc  
to  
jec  
gro  
par  
of

93

SOME REMARKS ON THE SYMPTOMS OF BRON-  
CHOCELE AND THE RESULTS OF OPER-  
ATIVE TREATMENT.<sup>1</sup>

By FRANCIS J. SHEPHERD, M.D.,

OF MONTREAL,

PROFESSOR OF ANATOMY AND LECTURER ON OPERATIVE SURGERY, M'GILL  
UNIVERSITY; SENIOR SURGEON TO THE MONTREAL  
GENERAL HOSPITAL.

SIMPLE goitre usually commences as a small kernel-like mass in one or other of the lobes of the thyroid and gradually increases in size as the years go on. Sometimes both sides of the gland are affected and also the isthmus, but in this country, at any rate, the unilateral variety is the most common. This form is usually encysted, and the cysts may be multiple or single, and may contain clear or blood-stained fluid, or the contents may be solid or semisolid colloid material. A cyst which has suddenly become larger from hæmorrhage very often, when it is tense, simulates a solid growth. In addition to these two forms of goitre we have cases where the gland is uniformly enlarged,—the interstitial or parenchymatous form. This kind is seen chiefly in young girls, especially at the time of puberty, and generally disappears in a few months; it increases markedly before the menstrual periods, and is often, when not of the soft, vascular variety, benefited by the administration of thyroid extract and iodides. In all these forms of goitre the general health is apt to be affected, the patients are more or less nervous, are subject to breathlessness on exertion, owing to pressure of the growth on the trachea and sometimes tachycardia; in fact, patients having the encysted solid forms are subjects of a kind of pseudo-Graves's disease, produced by the growth. The

<sup>1</sup> Address delivered before the Montreal Medico-Chirurgical Society, June, 1899.

relief afforded by removal of the growth is often very marked, as the following case illustrates:


Mrs. H., widow, aged thirty-five, a telegraph-operator, consulted me December, 1894; is a delicate, highly nervous woman, with some exophthalmos; has had enlargement of her neck since childhood. "Seven years ago the growth suddenly enlarged, but the enlargement disappeared temporarily on application of an ointment. This increase was accompanied by palpitation of the heart and great nervous excitability; from that time the gland slowly enlarged, until last summer, when it grew much more rapidly. Of late she has had frequent attacks of tachycardia and is very nervous, so much so that she had to give up her occupation. She has great difficulty in breathing, especially when she has to exert herself. In this case the left lobe is the larger, but both lobes extend from the hyoid bone to the clavicle. The growth seems to consist of a number of cysts, fluid and solid." Operation was performed, and the patient got rapidly well; all nervousness and tachycardia disappeared, her eyes became normal, and she was able to resume her work.


I have many such cases in my note-book. Here we have symptoms produced by increase of thyroid tissue of the nature of Graves's disease, and perhaps pressure also has something to do with it. I have quite recently operated on a case sent me by Dr. Birkett, where pressure from enlargement of the right lobe of the thyroid caused contraction of the pupil on that side and well-marked ptosis. Operation in this case is too recent to state results, but when I last saw her, ten days after the operation, the ptosis was certainly less marked.

The symptoms of Graves's disease are familiar; the highly nervous, excitable individual, with prominent eyes, enlarged, soft, vascular thyroid, rapid pulse, tremors, and often pyrexia,—symptoms which may be produced by taking thyroid internally. Such cases are exactly the opposite of myxœdema, due to absence or loss of thyroid tissue. Here the patient is dull, sleepy, has a stupid expression of face, low pulse and body temperature, dry, thick skin, with loss of hair. The patient sits about stupidly idle and sleeping most of the day. One disease is the exact opposite of the other, the for-



A case of solid colloid cyst, with symptoms of Graves's disease.





Colloid cyst after operation.

mer due to too much thyroid and the latter to too little. The exactly opposite conditions in these two diseases seem to me to prove the theory that too much thyroid is the cause of all that group of symptoms called exophthalmic goitre or Graves's disease, and in the cases I have seen of Graves's disease, which commenced with enlargement of the thyroid, the severity of the symptoms increased *pari passu* with the continued enlargement of the gland. This tends to prove the theory that increase of thyroid tissue is followed by symptoms of Graves's disease, and when the increased tissue is removed, the patient's health, in many cases, returns to normal, the symptoms of Graves's disease gradually disappearing. The following case illustrates this:

Miss L., aged twenty, sent me by Dr. Cornell, of Brockville, Ontario, first noticed enlargement of the thyroid some years ago. Both sides were enlarged and the gland continued enlarging. A short time after the enlargement, nervous symptoms developed, such as tachycardia and tremors. These increased in severity so that on the slightest exertion her pulse was almost uncountable. Then came exophthalmos, with persistent pyrexia, and oedema of the lower extremities; there was also great difficulty in breathing, especially on exertion. When I saw her, in 1896, she was a pale, anæmic girl, very thin, with exophthalmos and a rapid pulse,—140 to 200. She was excessively nervous, had tremors, some fever, and oedema of the lower extremities. Both lobes of the gland were considerably enlarged; the enlargement, whilst on both sides, was not the usual vascular, spongy enlargement of Graves's disease, but seemed to be made up of a separate solid cyst in each lobe. I advised operation, which was performed November 18, 1896. I enucleated from each lobe, by separate incisions, a solid cyst, the size of a small orange, full of colloid matter. The patient rapidly recovered from the operation, and was sent home in about ten days much better. I had a letter from her doctor May 18, 1899, in which he says: "I am pleased to tell you she is in excellent health; the respiratory trouble is of the past; the exophthalmos, the tachycardia, the anæmia and pyrexia are no more present, and she is perfectly well."

In other cases, although the symptoms may not be marked, the relief from operation is almost as great.

Jennie R., aged twenty-four, has had, since she was fifteen years old, enlargement of the thyroid. It commenced as a small, round growth in the right side, and gradually grew to its present size. For some years, owing to tachycardia and breathlessness, she has not been able to walk any distance or to go upstairs; for some years has had exophthalmos. When I saw her, early in January of this year, she was an anæmic girl with prominent eyes, a very nervous manner, and rapid pulse. She had a very large swelling, which was round and smooth, extending from the sternum to the hyoid bone, and this swelling went under the sterno-mastoid. It was not fluctuating, soft, and did not appear to be vascular. I looked upon the case as one of large colloid cyst, and recommended operation. This was performed January 27, 1899, and the tumor, as expected, turned out to be a colloid cyst. Hæmorrhage was quite free, and a number of ligatures had to be applied, the superior thyroid being ligated. Recovery was complicated by a continuous high temperature ( $104^{\circ}$  F.) and a very rapid pulse (180-200), following immediately on the operation. The discharges from the wound were tested and found perfectly sterile. It was supposed that this was a case of the so-called thyroid intoxication, which has been described by several writers. On giving free vent to the discharges from the wound, which were thin and watery, the temperature and pulse rapidly subsided, and the patient recovered completely. I heard from her on the 15th May. She says, "I have begun to feel like a different person, and just wonder how I ever put in such a miserable existence as I did the last five years. You would scarcely recognize me as the same person. My heart does not palpitate as it did before the operation."

I think these cases sufficiently prove that increase of thyroid tissue can produce a group of symptoms very much resembling Graves's disease. Now, these symptoms, in my experience, only come on in encysted cases with solid cysts. In cysts with fluid contents I have never seen them, so pressure can be but a small factor in the production of these symptoms.

In cases of true Graves's disease the improvement after operation is not so great. During the past four or five years I have operated on several cases, removing one-half of the enlarged thyroid. Although in these cases there has been

improvement, still it is not so rapid or so marked as in those cases where the disease in the gland is localized. In one of my recent cases, operated on in February last, the patient had all the chief symptoms of Graves's disease, such as tremors, tachycardia, pyrexia, etc., and a very large vascular thyroid. Operation relieved and her general health was much improved, but she writes me (May 23) that the nervousness still continues, though the tachycardia and exophthalmos are much better, and the remaining half of the gland is much smaller. In cases of true Graves's disease operation is not without danger. It seems that the danger is chiefly due to the anæsthetic; so much is this so that Kocher has given up general anæsthesia in these cases and resorts to local anæsthesia by cocaine. Even with local anæsthesia the operation is a dangerous one, and of Kocher's last fifteen cases of operation in Graves's disease two died.

It is my custom to advise operation in all rapidly growing goitres, especially if they be tumors of the solid form. If there be dyspnœa, the operation is urgently needed, but even if there be no dyspnœa, it is well to advise removal of those which are of recent formation and rapidly increasing in size, so that the serious train of symptoms which is characteristic of Graves's disease may be avoided.

*Operative Procedures.*—It is always well to be guided by the kind of case in choosing the form of operation. In the simple cystic case, where the cysts are large and not more than one or two in number, I invariably enucleate by the method I have described before (*ANNALS OF SURGERY*, Vol. xxii, p. 289), a simple incision over the cyst through skin and muscles down to the gland, tying the anterior jugular, if it be seen. When the gland is reached it is incised down to the bluish-white capsule of the cyst. The recognition of this capsule is most important, and when reached the cyst can be easily turned out. It has been my practice to open the cyst and evacuate its contents, so that it then can be pulled out of a small opening, and any vessels which bleed can be easily seized, as they are torn in separating the cyst. In some cases

of adherent cyst the separation is very difficult, but in fluid cysts the vascularity is not so great, nor is there so apt to be an adherent capsule. In the solid, colloid, encysted growths the enucleation is more difficult, owing to the greater vascularity. It is important here to get into the proper capsule, preferably the deeper one, for the superficial one is often covered with the ramifications of blood-vessels. Even in these cases before enucleating I open the tumor and remove some of its contents; where this is done the subsequent extraction is much less difficult. In diffuse cases or interstitial cases, and the true vascular thyroid of Graves's disease, in malignant disease, or where the cysts are multiple and small, or where the growth is very large, I prefer now to excise the gland. In Graves's and the interstitial cases only one lobe is excised. In these cases I make use of an incision down the inner border of the sterno-mastoid to near the upper border of the sternum, and then continue the incision transversely inward as far as necessary. Here the most important point is the free opening of the capsule of the gland. As soon as the capsule is divided the gland can be delivered, and the vessels tied without much difficulty. The superior thyroid artery should first be secured, then the gland thrown up and over to the opposite side. The inferior thyroid artery should be tied and not cut, and then the recurrent laryngeal nerve looked for and carefully separated. It runs up the posterior part of the gland; when the gland is enlarged it appears as if it entered it. The branches of the inferior thyroid artery with which it entwines should be cut near the gland, and also the veins which accompany these branches. I have cut the recurrent nerve once, and it was immediately sutured; the function has been partially recovered since.

*The After-Treatment.*—In the cases where enucleation has been performed there may be free oozing from the bed in which the cyst lay, and to prevent excessive oozing I pack this with strips of aseptic gauze, which I remove on the second day. In the cases where a portion of the gland has been removed a drain is inserted for twenty-four hours. The

wound is closed with horse-hair sutures and ordinary dry dressings applied. Usually the enucleation cases are discharged from hospital in a week or ten days.

The advantage of the enucleation method in suitable cases is the ease with which the operation is performed, the absence of risk of myxœdema, and the fact that the recurrent laryngeal nerve is never injured. The disadvantages are the chance of recurrence of the growth, and the tendency to oozing after operation; this oozing occasionally going on to secondary hæmorrhage. In nearly fifty cases of enucleation I have had recurrence in two,—in one on the opposite side, and in the other—a very small cyst—on the side that had previously been operated on. Both came to me, of their own accord, to have the cysts removed while they were small. Secondary hæmorrhage I have seen thrice,—once in a young man, who afterwards told me that he never had a tooth pulled without its bleeding for a week, and the other two were in women who had an apparent tendency to bleed. The blood oozed into the cavity which was left by the removal of the tumor, and only attracted notice when the breathing became difficult. In those cases the gauze had been removed too soon, for, after removal, on the second day there was a good deal of oozing. In such cases the gauze should have been replaced for another twenty-four hours.

Should secondary hæmorrhage occur, the wound should be laid open, the clots turned out, and the cavity firmly packed with sterile gauze. Swabbing out with tincture of the perchloride of iron, in the more severe cases, will arrest the hæmorrhage.

The cases I have seen, all recovered promptly, and the scar left did not appear to be any greater than that left after union by first intention.

I have seen many malignant cases, but have only operated on three. All subsequently died of recurrence of the disease in the lungs. Unless the tumor is removed very early there is little hope of permanent relief. In all the cases I have seen the gland had been enlarged for years before it took on a malignant action.